

Florpyrauxifen-benzyl Chemical Fact Sheet

Formulations

Florpyrauxifen-benzyl was registered with the EPA for aquatic use in 2017. The active ingredient is 2-pyridinecarboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoro-, phenyl methyl ester. The current Wisconsin-registered formulation is a liquid (ProcellaCOR™ EC) solely manufactured by SePRO Corporation.

Aquatic Use and Considerations

Florpyrauxifen-benzyl is a systemic herbicide that is taken up by aquatic plants. The herbicide is a member of a new class of synthetic auxins, the arylpicolinates, that differ in binding affinity compared to other currently registered synthetic auxins. The herbicide mimics the plant growth hormone auxin that causes excessive elongation of plant cells that ultimately kills the plant. Susceptible plants will show a mixture of atypical growth (larger, twisted leaves, stem elongation) and fragility of leaf and shoot tissue. Initial symptoms will be displayed within hours to a few days after treatment with plant death and decomposition occurring over 2 – 3 weeks. Florpyrauxifen-benzyl should be applied to plants that are actively growing; mature plants may require a higher concentration of herbicide and a longer contact time compared to smaller, less established plants.

Florpyrauxifen-benzyl has relatively short contact exposure time (CET) requirements (12 – 24 hours typically). The short CET may be advantageous for localized treatments of submersed aquatic plants, however, the target species efficacy compared to the size of the treatment area is not yet known.

In Wisconsin, florpyrauxifen-benzyl may be used to treat the invasive Eurasian watermilfoil (*Myriophyllum spicatum*) and hybrid Eurasian watermilfoil (*M. spicatum* X *M. sibiricum*). Other invasive species such as floating hearts

(*Nymphoides* spp.) are also susceptible. In other parts of the country, it is used as a selective, systemic mode of action for spot and partial treatment of the invasive plant hydrilla (*Hydrilla verticillata*). Desirable native species that may also be negatively affected include waterlily species (*Nymphaea* spp. and *Nuphar* spp.), pickerelweed (*Pontederia cordata*), and arrowhead (*Sagittaria* spp.).

It is important to note that repeated use of herbicides with the same mode of action can lead to herbicide-resistant plants, even in aquatic plants. Certain hybrid Eurasian watermilfoil genotypes have been documented to have reduced sensitivity to aquatic herbicides. In order to reduce the risk of developing resistant genotypes, avoid using the same type of herbicides year after year, and utilize effective, integrated pest management strategies as part of any long-term control program.

Post-Treatment Water Use Restrictions

There are no restrictions on swimming, eating fish from treated waterbodies, or using water for drinking water. There is no restriction on irrigation of turf. Before treated water can be used for non-agricultural irrigation besides turf (such as shoreline property use including irrigation of residential landscape plants and homeowner gardens, golf course irrigation, and non-residential property irrigation around business or industrial properties), follow precautionary waiting periods based on rate and scale of application, or monitor herbicide concentrations until below 2 ppb. For agricultural crop irrigation, use analytical monitoring to confirm dissipation before irrigating. The latest approved herbicide product label should be referenced relative to irrigation requirements.

Herbicide Degradation, Persistence and Trace Contaminants

Florpyrauxifen-benzyl is broken down quickly in the water by light (i.e., photolysis) and is also subject to microbial breakdown and hydrolysis. It has a half-life (the time it takes for half of the active ingredient to degrade) ranging from 1 – 6 days. Shallow clear-water lakes will lead to faster degradation than turbid, shaded, or deep lakes.

Florpyrauxifen-benzyl breaks down into five major degradation products. These materials are generally more persistent in water than the active herbicide (up to 3 week half-lives) but four of these are minor metabolites detected at less than 5% of applied active ingredient. EPA concluded no hazard concern for metabolites and/or degradates of florpyrauxifen-benzyl that may be found in drinking water, plants, and livestock.

Florpyrauxifen-benzyl binds tightly with surface sediments, so leaching into groundwater is unlikely. Degradation products are more mobile, but aquatic field dissipation studies showed minimal detection of these products in surface sediments.

Impacts on Fish and Other Aquatic Organisms

Toxicity tests conducted with rainbow trout, fathead minnow, water fleas (*Daphnia* sp.), amphipods (*Gammarus* sp.), and snails (*Lymnaea* sp.) indicate that florpyrauxifen-benzyl is not toxic for these species. EPA concluded florpyrauxifen-benzyl has no risk concerns for non-target wildlife and is considered "practically non-toxic" to bees, birds, reptiles, amphibians, and mammals.

Florpyrauxifen-benzyl does not bioaccumulate in fish or freshwater clams due to rapid metabolism and chemical depuration.



Human Health

EPA has identified no risks of concern to human health since no adverse acute or chronic effects, including a lack of carcinogenicity or mutagenicity, were observed in the submitted toxicological studies for florpyrauxifen-benzyl regardless of the route of exposure. EPA concluded with reasonable certainty that drinking water exposures to florpyrauxifen-benzyl do not pose a significant human health risk.

For Additional Information

Environmental Protection Agency Office of Pesticide Programs
www.epa.gov/pesticides

Wisconsin Department of Agriculture, Trade, and Consumer Protection
<http://datcp.wi.gov/Plants/Pesticides/>

Wisconsin Department of Natural Resources
608-266-2621
<http://dnr.wi.gov/lakes/plants/>

National Pesticide Information Center
1-800-858-7378
<http://npic.orst.edu/>

Washington State Department of Ecology. 2017.
<https://fortress.wa.gov/ecy/publications/documents/1710020.pdf>